



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# Effect of Aging on the Toughness of Human Cortical Bone: Evaluation by R-Curves

J.H. Kinney, R.K.Nalla, J.J.Kruzic, R.O.Ritchie

October 15, 2004

Bone

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

**Materials Sciences Division, Lawrence Berkeley National Laboratory, and  
Department of Materials Science and Engineering,  
University of California at Berkeley**

---

## **EFFECT OF AGING ON THE TOUGHNESS OF HUMAN CORTICAL BONE: EVALUATION BY R-CURVES**

---

***R. K. Nalla<sup>1</sup>, J. J. Kruzic<sup>1</sup>, J. H. Kinney<sup>2</sup>, and R. O. Ritchie<sup>1,3</sup>***

<sup>1</sup>Materials Sciences Division, Lawrence Berkeley National Laboratory, and  
Department of Materials Science and Engineering  
University of California, Berkeley CA 94720

<sup>2</sup>Lawrence Livermore National Laboratory, Livermore, CA 94550

<sup>3</sup>Corresponding author:  
Department of Materials Science and Engineering,  
381 Hearst Mining Building,  
University of California, Berkeley, CA 94720-1760  
Tel: (510) 486-5798; Fax: (510) 486-4881  
*E-mail address:* RORitchie@lbl.gov (R. O. Ritchie)

submitted March 2004  
revised June 2004

*Rapid Communications* submitted to *Bone*

This work was supported by the National Institutes of Health under Grant No. 5R01 DE015633 (for RKN), by the Director, Office of Science, Office of Basic Energy Science, Division of Materials Sciences and Engineering, Department of Energy under No. DE-AC03-76SF00098 (for JJK and ROR), and by the Laboratory Science and Technology Office, LLNL, under the auspices of the US Department of Energy W-7405-ENG48 (JHK).

# Effect of Aging on the Toughness of Human Cortical Bone: Evaluation by R-Curves

R. K. Nalla<sup>a</sup>, J. J. Kruzic<sup>a</sup>, J. H. Kinney<sup>b</sup>, and R. O. Ritchie<sup>a,\*</sup>

<sup>a</sup> Materials Sciences Division, Lawrence Berkeley National Laboratory, and  
Department of Materials Science and Engineering  
University of California, Berkeley CA 94720

<sup>b</sup> Lawrence Livermore National Laboratory, Livermore CA 94550

**Abstract:** Age-related deterioration of the fracture properties of bone, coupled with increased life expectancy, are responsible for increasing incidence of bone fracture in the elderly, and hence, an understanding of how its fracture properties degrade with age is essential. The present study describes *ex vivo* fracture experiments to quantitatively assess the effect of aging on the fracture toughness properties of human cortical bone in the longitudinal direction. Because cortical bone exhibits rising crack-growth resistance with crack extension, unlike most previous studies the toughness is evaluated in terms of resistance-curve (R-curve) behavior, measured for bone taken from wide range of age groups (34-99 years). Using this approach, both the *ex vivo* crack-initiation and crack-growth toughness are determined and are found to deteriorate with age; the initiation toughness decreases some 40% over six decades from 40 to 100 years, while the growth toughness is effectively eliminated over the same age range. The reduction in crack-growth toughness is considered to be associated primarily with a degradation in the degree of extrinsic toughening, in particular involving crack bridging in the wake of the crack.

**Keywords:** Cortical bone, aging, fracture toughness, R-curve.

---

\* Corresponding author: Tel: (510) 486-5798; Fax: (510) 486-4881  
E-mail address: RORitchie@lbl.gov (R. O. Ritchie)

## 1. Introduction

Aging related changes to the musculoskeletal system are known to increase the susceptibility of bone fracture (15), and in the case of the very elderly, consequent fractures can lead to mortality (16). Traditional thinking concerning “bone quality” has focused on bone mass or bone mineral density (BMD) as a predictor of such fracture risk. However, there is mounting evidence that BMD alone may not be the sole factor responsible for the aging-induced fracture risk (4, 14, 15). This has led to a renewed interest into how aging can alter the various mechanical properties of bone, and in particular the fracture resistance. Indeed, several studies that have looked at age-related issues in the mechanical properties of bone have shown a significant deterioration in the fracture toughness with age (2, 7-12, 29, 39, 40, 44-46). In order to characterize the deterioration of bone with age, most studies have utilized the fracture toughness,  $K_{Ic}$ , or the strain-energy release rate,  $G_{Ic}$ , as a single-parameter approach to characterize the resistance to fracture. However, in many materials, including cortical bone, so-called extrinsic toughening mechanisms<sup>1</sup>, such as constrained microcracking or crack bridging (24, 26), are active. For these materials, the fracture resistance increases with crack extension and stable crack growth can occur prior to unstable fracture. Although this necessitates a “resistance-curve” approach to evaluate the fracture toughness (20), such R-curves have only been utilized formally in relatively few studies (22, 25, 26, 28, 35-38, 41) to characterize bone fracture.

---

<sup>1</sup> Crack propagation can be considered as a mutual competition between two classes of mechanisms: *intrinsic* mechanisms that operate ahead of the crack tip, and affect the material’s inherent resistance to fracture and damage, and *extrinsic* mechanisms that principally operate in the wake of the crack tip, and “shield” the crack from the applied driving force (13, 30, 31). Whereas intrinsic mechanisms primarily govern the crack-initiation toughness, extrinsic mechanisms, specifically crack bridging in bone (25), operate in the crack wake and govern the crack-growth toughness. As the effect of extrinsic mechanisms is dependent on the size of the crack, this leads to rising R-curve behavior.

An early application of the R-curve approach for bone was presented by Vashishth *et al.* (36) for studies on cracking in human and bovine tibia in the longitudinal (anatomically proximal-distal) orientation. They reported rising R-curves for both cases, with steeper curves in bovine samples, reflecting the higher crack-growth toughness of the bovine bone. They later reported similar R-curve toughening in red deer antler (38). Another study, by Pezzotti and Sakakura, reported a rising R-curve in bovine bone (specimen orientation unclear) (28); however after an initial rising portion, a steady-state “plateau” toughness was achieved, as seen in a number of materials that exhibit R-curve behavior (19, 20). Malik *et al.* (22) reported rising R-curve behavior which displayed such a plateau (and in some cases subsequently decreased) for transverse crack growth in equine bone. The most recent work on R-curve behavior in human bone, by the present authors, investigated *ex vivo* longitudinal (proximal-distal) crack growth in the humerus; linear rising R-curves were also observed in that study (25, 26). In addition, this work provided definitive evidence that crack bridging by uncracked ligaments, rather than a mechanism involving constrained microcracking which had been previously suggested (36-38), is the dominant extrinsic toughening mechanism responsible for the toughening and rising R-curve behavior in bone.

While it is now well established that the fracture toughness of bone must be characterized in terms of rising R-curves, there is a paucity of relevant data on how such R-curve behavior changes with aging. Very limited data (only five total R-curves) on the R-curve behavior of human femoral bone demonstrated a decrease in both initiation and growth toughness with age (41), although no mention of mechanisms was made. Accordingly, in the present paper, we seek to investigate the *ex vivo* R-curve fracture

properties of bone as a function of age in order to 1) provide additional data to confirm this aging effect and 2) to elucidate the mechanisms responsible for any age-related changes.

## 2. Materials and Methods

Fresh frozen human cadaveric humeral cortical bone was used in this study from nine donors. The age of these donors varied from 34 to 99 years old (cause of donor death unrelated to skeletal state); the gender of the donors together with anatomical location are given in Table I. Blocks of bone were obtained by carefully sectioning the medial cortices of the mid-diaphyses of the humeri. Seventeen ( $N = 17$ ) compact-tension, C(T), specimens, with specimen thicknesses,  $B \sim 1.2\text{-}3.3\text{ mm}$ <sup>2</sup>, widths,  $W \sim 13\text{-}18.3\text{ mm}$  and initial crack lengths,  $a \sim 3.0\text{-}5.5\text{ mm}$ , were machined from these blocks, and divided into three age groups - arbitrarily named *Young*, *Middle-Aged* and *Aged* (see Table I for details). The samples were all orientated with the starter notch and the nominal crack-growth direction along the proximal-distal direction of the humerus (in the longitudinal-radial plane), i.e., parallel to the long axis of the osteons and hence, long axis of the humerus. This orientation is designated C-L according to ASTM Standard E399 for fracture toughness testing (1), where the first letter of the designation refers to the direction normal to the crack plane (circumferential) and the second refers to the cracking direction (longitudinal). Although based on expected physiological loading conditions, the transverse, and not longitudinal, cracking direction, would seem to be the most relevant, cracks loaded so as to cause growth in the transverse direction *ex vivo* have been found to deflect towards the longitudinal direction (e.g., (6)). Accordingly, in order to gain a mechanistic understanding of cortical bone fracture, the longitudinal orientation would appear to be physiologically

relevant, and the larger amount of crack extension that is possible in this orientation is useful experimentally.

R-curves were measured to evaluate the resistance to fracture in terms of the stress intensity,  $K$ , as a function of crack extension,  $\Delta a$ , under a monotonically increasing driving force. The C(T) specimens were thawed and thoroughly hydrated prior to testing by soaking in Hanks' Balanced Salt Solution (HBSS) for at least 40 hr at room temperature in air-tight containers. Tests were then conducted in ambient air (25°C, 20-40% relative humidity) with the specimens being continuously irrigated with HBSS. The specimens were loaded in displacement control using standard servo-hydraulic testing machines (MTS 810, MTS Systems Corporation, Eden Prairie, MN) with a loading rate  $\sim 0.015$  mm/s until the onset of cracking, which was determined by a drop in load, or non-linearity in the load-displacement curve. At this point, the sample was manually unloaded by 10-20% of the peak load to record the sample load-line compliance at the new crack length using a linear variable-displacement transducer (LVDT) mounted in the load frame. This process was repeated at regular intervals until the end of the test (arbitrarily chosen when data for at least 4 mm of crack growth was obtained), at which point the compliance and loading data were analyzed to determine fracture resistance,  $K_R$ , as a function of crack extension,  $\Delta a$ . Crack lengths,  $a$ , were calculated from the compliance data obtained during the test using standard C(T) load-line compliance calibrations (32).<sup>3</sup> Further details of the testing procedures are provided elsewhere (25, 26). Statistical analysis of the data was conducted

---

<sup>2</sup> Variations in sample thickness were related to variations in the cortex thickness for the various donors. They are not expected to influence the measured toughness; indeed Norman *et al.* reported no effect of specimen thickness on the toughness of human bone C(T) specimens of a 2 – 3 mm range (27).

<sup>3</sup> The use of elastic compliance is a widely used technique in fracture mechanics testing to accurately measure crack length. The calibrations which relate compliance to crack length are valid for isotropic and anisotropic linear-elastic solids (32) and have been utilized successfully for numerous materials including dentin and bone (25, 26).



using the non-parametric Kruskal-Wallis test. The data were also subjected to linear regression analysis against age.

To observe crack-microstructure interactions, synchrotron x-ray computed tomography (SRCT) was performed on two specimens of the *Young* (data reported in ref. (25)) and *Aged* groups. This work was performed in part at the Stanford Synchrotron Radiation Laboratory (SSRL), Menlo Park, CA, and at the Advanced Light Source (ALS), Berkeley, CA. Imaging was performed with monochromatic x-rays (25 keV at SSRL and 18 keV at ALS), with a voxel size (spatial resolution) of  $\sim 5 \mu\text{m}$ . The tomography data were reconstructed using a Fourier-filtered back-projection algorithm; further details of this technique are described elsewhere (17, 18).

### 3. Results

As described above, the *ex vivo* load-displacement data obtained were analyzed to evaluate the resistance to fracture in terms of the stress intensity,  $K$ , as a function of crack extension,  $\Delta a$ . The resulting monotonically rising R-curves for hydrated cortical bone are shown in Fig. 1. The *crack-initiation toughness*,  $K_0$ , was obtained by extrapolating a linear fit of the data for each sample to  $\Delta a = 0$ , while the (linear) slope of the R-curve gave a measure of the *crack-growth toughness*. Mean (non-weighted)  $K_0$  values of 2.07 (S.D. = 0.11), 1.96 (S.D. = 0.15), and 1.26 (S.D. = 0.22)  $\text{MPa}\sqrt{\text{m}}$ , and mean slopes of 0.37 (S.D. = 0.06), 0.16 (S.D. = 0.01), and 0.06 (S.D. = 0.04)  $\text{MPa}\sqrt{\text{m/mm}}$  were thus obtained for the *Young*, *Middle-Aged*, and *Aged* groups, respectively; individual values are listed in Table I. Statistical analysis using the Kruskal-Wallis test of the data indicated that, for the three age groups, variation among group medians was significant ( $p = 0.025$  and  $0.0036$  for the initiation and the growth toughness, respectively). *Post-hoc* analysis was not performed in

view of the small sample sizes. The data in Table I are plotted in Fig. 2 as variations of the crack-initiation and growth toughnesses as a function of age. While the initiation toughness decreases with age, the effect of aging is more evident on the growth toughness which is essentially eliminated in the *Aged* group.

Figs. 3a-b shows typical two-dimensional through-thickness “slices” obtained by SRCT for specimens belonging to the *Young* and *Aged* groups at various distances behind the crack tip. There is evidence of “crack bridging” in both cases from the formation of uncracked ligaments in the crack wake; these are intact regions, often tens of micrometers in size (i.e., substantially larger than individual collagen fibers), that form along the crack path, either by the non-uniform advance of the crack front and/or by the imperfect linking of microcracks that initiated ahead of the crack tip with the main crack. It has been previously shown that such bridging is the primary mechanism of toughening for cracking in the longitudinal orientation in bone and is responsible for the rising R-curve behavior (25, 26). Further examination of the tomographic slices revealed that there is a definitive decrease in the size and number of these bridges with age. Fig. 3c shows the variation in the area fraction of such bridges with distance from the crack tip for *Young* and *Aged* bone. It is apparent that the bridging zones are larger in *Young* bone (roughly 5.5 mm vs. 3.5 mm), and within the zone, the area fractions are in general larger for that group. Such observations are consistent with the reduction in the growth toughness with age, as discussed below.

#### **4. Discussion**

The data in Table I are plotted in Fig. 2 as variations of the crack-initiation and growth toughnesses as a function of age. The only other R-curve data available for human

cortical bone from Vashishth *et al.* (36) and from Wu and Vashishth (41) are also included. While the initiation toughness data from these studies agrees well with the trend suggested by the linear regressions in Fig. 2a, there is a stronger effect of age on the growth toughness in ref. (41) as compared to the present study. It should be noted that the bone used in those studies were from a different anatomical location (tibia in ref. (36) and femur in ref. (41)). There is a clear, observable trend of decreasing toughness with age in the present study; specifically, the crack-initiation toughness decreases by ~40% over six decades from 40 to 100 years, while the growth toughness is essentially eliminated over the same age range. Such deterioration in the fracture resistance with age is consistent with the trend observed in studies that report single-value toughnesses (e.g., (2, 7-12, 29, 39, 40, 44-46)). What is important about these results, together with those of Wu and Vashishth (41), is that they clearly show that not only the intrinsic resistance to fracture (as reflected by the crack-initiation toughness), but also the increasing resistance to crack propagation (as reflected by the crack-growth toughness) decreases with age (Figs. 1,2b). Indeed, the age-related deterioration in the crack-growth toughness is clearly the more dominant effect.

As noted above, the crack-growth toughness is reflective of the contribution from extrinsic toughening mechanisms, which in bone are principally associated with crack bridging (25, 26). The prime source of such bridging in human bone appears to be from the formation of uncracked ligaments in the crack wake (Fig. 3). The results in Fig. 2b strongly imply that the contribution from such a mechanism is markedly reduced with age. The magnitude of this contribution is dictated by the size of the bridging zone, the area fraction of the bridges in the zone, and their load-bearing capacity. All of these factors

may change with aging; indeed, Fig. 3 shows that there is a lower density of such bridges in older bone. A more detailed quantitative assessment of these parameters is currently being undertaken.

Many effects of aging on bone have been studied, in particular in association with the micro/ultra-structural changes (i.e., at micron- and nano-scale dimensions) induced by a number of parameters, including increased mineralization (3), increased microdamage (33), lowered collagen quality (39), and increased bone turnover (23). However, the underlying mechanisms that result in age-related changes in the fracture resistance are poorly understood. The increased mineralization has been implicated in reduced elastic deformability (2), and the (single-value) fracture toughness (12). Increased levels of microdamage (microcracking) have been shown to lower the fracture resistance (42), presumably by lowering the intrinsic toughness. Possible changes in collagen network integrity with age (39) could result in weaker bridges, and hence, a lower growth toughness in older bone. More specifically, age is known to increase non-enzymatic cross-linking in the collagen (39); as this reduces the post-yield deformation of the collagen, this also has been used to explain the age-induced reduction in growth toughness via a microcracking model (36), although recent studies have cast doubt on the significance of microcracking in affecting the toughness of bone (25, 26). Finally, it has been suggested that elevated bone turnover in older bone (21), though beneficial in repairing damage, may have a deleterious effect on the toughness due to the formation of resorption cavities and hence increased porosity. Elevated turnover also results in a higher density of secondary osteons (23), and associated cement lines which are known to provide weak interfaces, and hence preferred (weaker) paths, for cracking (5, 24, 25, 34, 43). In fact, a recent study has

concluded that elevated turnover is a risk factor in its own right, independent of its actions on BMD (14).

Thus, although the age-induced decrease in the fracture toughness of bone has been clearly quantified, specifically in terms of resistance to both crack initiation and more importantly crack growth, the mechanistic reasons for this deterioration are as yet unclear. As the microstructural factors affecting crack initiation and growth in most materials are invariably quite distinct (13, 30, 31), the challenge is to identify *and quantify* the specific mechanisms affecting each process in terms of the changes that occur in the micro/ultra-structure of bone with age. Furthermore, other factors (e.g., gender, race, pathology, genetic reasons, etc.) that might have an effect on the micro/ultra-structure and consequently, the fracture behavior also need to be studied.

In summary, the fracture toughness of cortical bone, expressed in terms of a rising R-curve, shows significant deterioration with aging. In quantitative terms, the *ex vivo* crack-initiation toughness was reduced by ~40%, whereas the crack-growth toughness was effectively eliminated, as age increased from 34 to 99 years. These results demonstrate the need to interpret this deterioration in bone quality in terms of specific age-related changes in the micro/ultra-structure of bone that separately affect the crack initiation and growth stages of fracture, both for the purpose of fully characterizing the aging properties of bone and for identifying the actual mechanisms that ultimately increase its fragility and fracture susceptibility. In this regard, the present results have clearly identified that a primary mechanistic factor in the deterioration in the toughness of bone with age can be associated with a degradation in the crack bridging that is developed in the crack wake.

## **Acknowledgements**

This work was supported by the National Institutes of Health under Grant No. 5R01 DE015633 (for RKN), by the Director, Office of Science, Office of Basic Energy Science, Division of Materials Sciences and Engineering, Department of Energy under No. DE-AC03-76SF00098 (for JJK and ROR), and by the Laboratory Science and Technology Office, LLNL, under the auspices of the US Department of Energy W-7405-ENG48 (JHK). We acknowledge the support of X-ray wiggler beamline (BL 10-2) at the Stanford Synchrotron Radiation Laboratory (SSRL), supported by Department of Energy contract No. DE-AC03-76SF00515, and the dedicated tomography beamline (BL 8.3.2) at the Advanced Light Source (ALS), supported by the Department of Energy (BES) under Contract No. DE-AC03-76SF00098. Finally, we also wish to thank Prof. A. P. Tomsia for his support, and Drs. C. Puttlitz and Z. Xu for supply of the cortical bone.

## References

1. ASTM E399-90 (Reapproved 1997). Annual Book of ASTM Standards, Vol. 03.01: Metals- Mechanical Testing; Elevated and Low-temperature Tests; Metallography: ASTM, West Conshohocken, Pennsylvania, USA; 2002.
2. Akkus, O., Adar, F., and Schaffler, M. B. Age-related changes in physicochemical properties of mineral crystals are related to impaired mechanical function of cortical bone. *Bone* 34:443-453; 2004.
3. Akkus, O., Polyakova-Akkus, A., Adar, F., and Schaffler, M. B. Aging of microstructural compartments in human compact bone. *J Bone Miner Res* 18:1012-1019; 2003.
4. Aspray, T. J., Prentice, A., Cole, T. J., Sawo, Y., Reeve, J., and Francis, R. M. Low bone mineral content is common but osteoporotic fractures are rare in elderly rural Gambian women. *J Bone Miner Res* 11:1019-1025; 1996.
5. Behiri, J. C., and Bonfield, W. Orientation dependence of the fracture mechanics of cortical bone. *J Biomech* 22:863-872; 1989.
6. Behiri, J. C., and Bonfield, W. Orientation dependence on fracture mechanics of bone. *J Biomech* 22:863-872; 1989.
7. Bonfield, W., Behiri, J. C., and Charalamides, C. Orientation and age-related dependence of the fracture toughness of cortical bone. In: E. Schneider (ed.), *Biomechanics: Current Interdisciplinary Research*: Martinus Nijhoff Publishers, Dordrecht; 1985.
8. Brown, C. U., and Norman, T. L. Fracture toughness of human cortical bone from the proximal femur. *Advances in Bioengineering* 31:121-122; 1995.
9. Brown, C. U., Yeni, Y. N., and Norman, T. L. Fracture toughness is dependent on bone location- A study of the femoral neck, femoral shaft, and the tibial shaft. *J Biomed Mater Res* 49:380-389; 2000.
10. Burstein, A., Reilly, D., and Martens, M. Aging of bone tissue mechanical properties. *J Bone Joint Surgery* 58A:82-86; 1976.
11. Currey, J. D. Changes in impact energy absorption with age. *J Biomech* 12:459-469; 1979.
12. Currey, J. D., Brear, K., and Zioupos, P. The effects of ageing and changes in mineral content in degrading the toughness of human femora. *J Biomech* 29:257-260; 1996.
13. Evans, A. G. Perspective on the development of high toughness ceramics. *J Am Ceramic Soc* 73:187-206; 1990.
14. Heaney, R. Is the paradigm shifting? *Bone* 33:457-465; 2003.
15. Hui, S. L., Slemenda, C. W., and Johnston, C. C. Age and bone mass as predictors of fracture in a prospective study. *J Clin Invest* 81:1804-1809; 1988.
16. Jennings, A. G., and de Boer, P. Should we operate on nonagenarians with hip fractures? *Injury* 30:169-172; 1999.
17. Kinney, J. H., Haupt, D. L., Nichols, M. C., Breunig, T. M., Marshall, G. W., and Marshall, S. J. The X-Ray Tomographic Microscope - 3-Dimensional Perspectives of Evolving Microstructures. *Nuclear Instruments & Methods in Physics Research Section A-Accelerators Spectrometers Detectors and Associated Equipment* 347, pp. 480-486; 1994.

18. Kinney, J. H., and Nichols, M. C. X-Ray Tomographic Microscopy (XTM) Using Synchrotron Radiation. *Ann Review Mater Sci* 22, pp. 121-152; 1992.
19. Knott, J. F. *Fundamentals of fracture mechanics*. London, U.K.: Butterworth & Co. (Publishers) Ltd; 1976.
20. Lawn, B. R. *Physics of Fracture*. *J Am Ceramic Soc* 66:83; 1983.
21. Lee, T. C., Staines, A., and Taylor, D. Bone adaptation to load: Microdamage as a stimulus for bone remodelling. *J Anat* 201:437-446; 2002.
22. Malik, C. L., Stover, S. M., Martin, R. B., and Gibeling, J. C. Equine cortical bone exhibits rising R-curve fracture mechanics. *J Biomech* 36:191-198; 2003.
23. McCalden, R. W., McGeough, J. A., Barker, M. B., and Court-Brown, C. M. Age-related changes in the tensile properties of cortical bone. The relative importance of changes in porosity, mineralization, and microstructure. *J Bone Joint Surg Am* 75:1193-1205; 1993.
24. Nalla, R. K., Kinney, J. H., and Ritchie, R. O. Mechanistic fracture criteria for the failure of human cortical bone. *Nature Materials* 2:164-168; 2003.
25. Nalla, R. K., Kruzic, J. J., Kinney, J. H., and Ritchie, R. O. Mechanistic aspects of fracture and R-curve behavior in human cortical bone. *Biomaterials* 26:217-231; 2005.
26. Nalla, R. K., Kruzic, J. J., and Ritchie, R. O. On the Origin of the Toughness of Mineralized Tissue: Microcracking or Crack Bridging? *Bone* 34:790-798; 2004.
27. Norman, T. L., Vashishth, D., and Burr, D. B. Fracture toughness of human bone under tension. *J Biomech* 28:309-320; 1995.
28. Pezzotti, G., and Sakakura, S. Study of the toughening mechanisms in bone and biomimetic hydroxyapatite materials using Raman microprobe spectroscopy. *J Biomed Mater Res* 65A:229-236; 2003.
29. Phelps, J. B., Hubbard, G. B., Wang, X., and Agrawal, C. M. Microstructural heterogeneity and the fracture toughness of bone. *J Biomed Mater Res* 51:735-741; 2000.
30. Ritchie, R. O. Mechanisms of fatigue crack propagation in metals, ceramics and composites: Role of crack-tip shielding. *Mater Sci Engg* 103:15-28; 1988.
31. Ritchie, R. O. Mechanisms of fatigue-crack propagation in ductile and brittle solids. *Int J Fracture* 100:55-83; 1999.
32. Saxena, A., and Hudak Jr., S. J. Review and extension of compliance information for common crack growth specimens. *Int J Fracture* 14:453-468; 1978.
33. Schaffler, M. B., Choi, K., and Milgrom, C. Aging and matrix microdamage accumulation in human compact bone. *Bone* 17:521-525; 1995.
34. Taylor, D. Failure processes in hard and soft tissues. In: I. Milne, R. O. Ritchie, and B. L. Karihaloo (eds.), *Comprehensive Structural Integrity: Fracture of Materials from Nano to Macro* 9, pp. 35-96. Oxford, U.K.: Elsevier Inc.; 2003.
35. Vashishth, D. Rising crack-growth-resistance behavior in cortical bone: Implication for toughness measurements. *J Biomech* 37:943-946; 2004.
36. Vashishth, D., Behiri, J. C., and Bonfield, W. Crack growth resistance in cortical bone: Concept of microcrack toughening. *J Biomech* 30:763-769; 1997.
37. Vashishth, D., Tanner, K. E., and Bonfield, W. Contribution, development and morphology of microcracking in cortical bone during crack propagation. *J Biomech* 33:1169-1174; 2000.



38. Vashishth, D., Tanner, K. E., and Bonfield, W. Experimental validation of a microcracking-based toughening mechanism for cortical bone. *J Biomech* 36:121-124; 2003.
39. Wang, X., Shen, X., Li, X., and Agrawal, C. M. Age-related changes in the collagen network and toughness of bone. *Bone* 31:1-7; 2002.
40. Wang, X. D., Masilamani, N. S., Mabrey, J. D., Alder, M. E., and Agrawal, C. M. Changes in the fracture toughness of bone may not be reflected in its mineral density, porosity, and tensile properties. *Bone* 23:67-72; 1998.
41. Wu, P.-C., and Vashishth, D. Age related changes in cortical bone toughness: Initiation vs. propagation. 2nd Joint EMBS/BMES Conference 1, pp. 425-426. Houston, TX: IEEE; 2002.
42. Yeni, Y. N., and Fyhrie, D. P. Fatigue damage-fracture mechanics interaction in cortical bone. *Bone* 30:509-514; 2002.
43. Yeni, Y. N., and Norman, T. L. Calculation of porosity and osteonal cement line effects on the effective fracture toughness of cortical bone in longitudinal crack growth. *J Biomed Mater Res* 51:504-509; 2000.
44. Yeni, Y. N., and Norman, T. L. Fracture toughness of human femoral neck: Effect of microstructure, composition, and age. *Bone* 26:499-504; 2000.
45. Zioupos, P., and Currey, J. D. Changes in the stiffness, strength, and toughness of human cortical bone with age. *Bone* 22:57-66; 1998.
46. Zioupos, P., Currey, J. D., and Hamer, A. J. The role of collagen in the declining mechanical properties of aging human cortical bone. *J Biomed Mater Res* 2:108-116; 1999.

## List of Figure Captions

**Fig. 1:**  $K_R(\Delta a)$  resistance-curves for stable *ex vivo* crack extension in human cortical bone as a function of age. Note the linearly rising R-curve behavior. The inset schematically shows the anatomical orientation that the specimens were taken from the humeri.


**Fig. 2:** Variation in the (a) crack-initiation toughness ( $K_o$ ), and (b) crack-growth toughness (slope of the R-curve) with age for human cortical bone. A linear regression of the data is shown in each case (fit equation and coefficient of determination,  $R^2$ , is also included). Data from Vashishth *et al.* (36) and Wu and Vashishth (41) are also plotted for comparison (not included in regression).

**Fig. 3:** Two-dimensional computed x-ray tomographic reconstruction slices showing typical cracks in specimens taken from the (a) *Young* (34FL), and (b) *Aged* (85FR) groups. The numbers on top of each figure indicate the distance from the (nominal) crack tip, and the white arrows indicate uncracked-ligament “crack bridges”. (c) The fraction of such bridges with distance from the crack tip indicating smaller area fractions and bridging-zone size in the older bone.


**Table I:** Resistance-Curve Behavior of Human Cortical Bone with Age

Donor Information*	Initiation Toughness $K_0$ (MPa $\sqrt{m}$ )	Slope (MPa $\sqrt{m/mm}$ )	Coefficient of Determination ( $R^2$ )
<b>Young (34-41 years):</b>			
34FL	2.12	0.31	0.98
37ML	1.69	0.50	0.88
37ML	2.20	0.28	0.97
37MR	2.07	0.49	0.97
37MR	1.85	0.41	0.94
41FL	2.07	0.41	0.97
41FL	2.23	0.34	0.96
<b>Middle-Aged (61-69 years):</b>			
61ML	2.00	0.16	0.80
69FLa	2.09	0.16	0.49
69FLa	2.09	0.18	0.37
69FRb	1.66	0.23	0.91
69FRb	1.94	0.06	0.60
<b>Aged (85-99 years):</b>			
85FRa	1.44	0.02	0.40
85FRb	1.32	0.07	0.56
85FRb	1.30	0.08	0.73
99MR	1.11	0.07	0.65
99MR	0.92	0.11	0.47


\* The notation reads as follows: Age (years), Sex (M=Male, F=Female), Arm (L=Left, R=Right), with any subsequent letter being a unique identifier when more than one donor was of the same age (69 and 85 years). Data for 34-41 year group from ref. (25).



**Fig. 1:**  $K_R(\Delta a)$  resistance-curves for stable *ex vivo* crack extension in human cortical bone as a function of age. Note the linearly rising R-curve behavior. The inset schematically shows the anatomical orientation that the specimens were taken from the humeri.






(a)



(b)

**Fig. 2:** Variation in the (a) crack-initiation toughness ( $K_0$ ), and (b) crack-growth toughness (slope of the R-curve) with age for human cortical bone. A linear regression of the data is shown in each case (fit equation and coefficient of determination,  $R^2$ , is also included). Data from Vashishth *et al.* (36) and Wu and Vashishth (41) are also plotted for comparison (not included in regression).





(c)

**Fig. 3:** Two-dimensional computed x-ray tomographic reconstruction slices showing typical cracks in specimens taken from the (a) *Young* (34FL), and (b) *Aged* (85FR) groups. The numbers on top of each figure indicate the distance from the (nominal) crack tip, and the white arrows indicate uncracked-ligament “crack bridges”. (c) The fraction of such bridges with distance from the crack tip indicating smaller area fractions and bridging-zone size in the older bone.